


Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Archives of Psychiatric Nursing

journal homepage: www.elsevier.com/locate/apnu

Emotional intelligence of nurses caring for COVID-19 patients: A cross-sectional study

Seyed Tayeb Moradian^a, Mahmoud Movahedi^b, Mohammad Goudarzi Rad^c, Yaser Saeid^{d,*}

^a Atherosclerosis Research Center and Faculty of Nursing, Baqiyatallah University of Medical Sciences, Tehran, Iran

^b Department of Psychiatry Sciences Research Center, Faculty of Nursing, Baqiyatallah University of Medical Sciences, Tehran, Iran

^c Forghani Hospital, Qom University of Medical Sciences, Qom, Iran

^d Trauma Research Center and Faculty of Nursing, Baqiyatallah University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Keywords:
COVID-19
Emotional intelligence
Nurse

ABSTRACT

Purpose: The aim of this study was to assess the emotional intelligence of nurses caring for COVID-19 patients.
Methods: This was a descriptive cross-sectional study that was conducted from May to July 2020 in Tehran, Iran. Nurses caring for patients with COVID-19 were requested to fill in Bradbury and Graves's questionnaire online using a questionnaire in electronic format.

Results: Finally 211 nurses completed the questionnaires. Most of the nurses were working in critical care wards and caring for critical patients (61.6). Nurses' emotional intelligence was reported to be 63.19 (8.22). In general, the nurses' emotional intelligence was moderate. Between the dimensions, self-awareness and self-management had the highest scores. Also, the lowest score was related to self-management. The ward type and complexity of care had no effect on the scores of emotional intelligence. Nurses caring for patients with moderate disease severity had a higher relationship management score than nurses caring for critically ill patients ($P < 0.05$).

Conclusion: The total score of emotional intelligence was moderate. Due to the continuation of the COVID-19 pandemic and the possibility of mental and physical fatigue of health care workers, improving emotional intelligence can be effective in resilience and stability of the psychological status of employees.

Introduction

World health organization declared the COVID-19 crisis as a pandemic with a public health emergency (Priesemann et al., 2021; Smith et al., 2020). Health care providers have a high tendency to be infected with the virus. The prevalence of COVID-19 in health care workers (HCW) is more than the general population. HCWs are prone to psychological problems (Huang et al., 2020). Studies report that this pandemic could increase the risk of psychological symptoms in asymptomatic populations and exacerbate the symptoms in previously diagnosed psychological problems (Sommer et al., 2016). Anger, confusion, and post-traumatic stress disorder are reported as a consequence of this pandemic (Mo et al., 2020).

HCWs who are working in the frontline are at risk for psychological disorders and sometimes the prevalence of disorders such as depression and anxiety in HCWs is reported more than the general population (Montes-Berges & Augusto, 2007). As a result, we should take measures to improve the psychological health of frontline nurses to prevent

further damage to the nurses and their patients (Doherty et al., 2013). Emotional intelligence (EI) is one of the most important aspects that improve psychological health. EI is the ability to recognize, understand, and regulate emotions and use them in life (Morris & Wakefield, 2008). Also, EI is recognized as a core variable affecting job performance (Nightingale et al., 2018). It is also a key factor for Creating, reinforcing the emotional capacity that nurses need to interact with patients, and bearing the emotional burden (Morris & Wakefield, 2008). The level of EI may vary between different occupations, especially among health care professionals (Talman et al., 2020). Some studies report a strong correlation between EI and nurses' performance. They also report the EI of nurses high (Alonazi, 2020).

The intensifying of the COVID-19 pandemic plays a strong role in emotional and psychological disorders between HCWs especially nurses (Wu et al., 2020). The level of emotional intelligence during the COVID-19 crisis has been evaluated in previous studies (Alonazi, 2020; Baba, 2020). Improving emotional intelligence skills affects nurses' performance (Alonazi, 2020) and less experiencing negative emotions (Moroi

* Corresponding author.

E-mail address: yasernurse84@yahoo.com (Y. Saeid).

<https://doi.org/10.1016/j.apnu.2021.10.011>

Received 31 January 2021; Received in revised form 24 August 2021; Accepted 29 October 2021

Available online 6 November 2021

0883-9417/© 2021 Elsevier Inc. All rights reserved.

et al., 2021). This study was to assess the EI in nurses caring for COVID-19 patients. The results of present study besides having theoretical implications could open pathways for conducting more research.

Methods

Aim

This descriptive cross-sectional study was done from May to July 2020 in Tehran, Iran. The aim of the study was to assess the EI of nurses caring for COVID-19 patients.

Sample

All nurses who were working in COVID-19 wards were invited to participate in the study. Those with at least an associate degree in nursing who were caring for COVID-19 patients and had willing to participate in the study were included. Totally 211 nurses participated in the study.

Data collection

Sampling was done using a web-based online questionnaire. The questionnaire was sent online to 250 people via email and social media. The required time for responding the questionnaire was 5 to 7 min. If no response was given, after one week a reminder was sent. Finally, 211 questionnaires were completed. These questionnaires were assessed and analyzed using statistical software. Totally the response rate was reported 87.2%. The questionnaire had two parts. The first section was demographic and job description and the second part was the Emotional Intelligence questionnaire developed by Brad berry and Greaves in 2005.

All nurses included in the study worked in the wards where COVID-19 patients were admitted. Nurses from both private and governmental sectors were included. Also, they were working in both general and intensive care wards. Participants were informed about the study objective and the confidentiality of the data. They had the right to withdraw the study based on their desire. The informed written consent was taken from all participants.

This questionnaire has 28 items and four aspects of Self-awareness (questions 1–6), self-management (questions 7–15), social awareness (questions 16–21), and relationship management (questions 22–28). The scoring of this questionnaire is based on a Likert scale from 1 to 6. The sum of the score is the total score of each aspect. Then we computed them and calculate the total score of the questionnaire. For better understanding, the total score of the questionnaire was computed to 0 to 100. Having a higher score means higher EI. This questionnaire is translated and transcultural adapted in Persian. The reliability of this test in studies based on internal consistency method, Cronbach's alpha coefficient has been reported 0.83. The validity of this scale has been determined by a correlation coefficient of $r = 0.67$ and $P < 0.01$ (Ganji et al., 2006).

Ethical consideration

The present study was approved by the Ethics Committee of Baqiyatallah University of Medical Sciences, Tehran, Iran, with code IR.BMSU.REC.1399.227.

Statistical analysis

Data were analyzed using the Statistical Package for the Social Sciences (SPSS) 25.0 statistical package (Chicago, IL, USA), and a two-sided P -value > 0.05 was considered as a statistically significant difference. Categorical variables are presented as frequency rates and percentages, and continuous variables are described using mean \pm standard deviation

(SD) values. Data analysis was performed using descriptive and inferential statistics (independent t -test, Multinomial logistic regression, Spearman's rho correlation coefficient). The normality of numeric variables was tested using the Kolmogorov-Smirnov test. The comparison of demographic characteristics and job experience parameters was done by independent t -test for continuous variables and Chi-square test or Fisher's exact test for categorical variables.

Results

Totally 211 nurses participated in the study. Of these, 52.7% of them were female, 71.6% were married, and 72.5% of them were BSc. The mean age and work experience of participants were 34 (7.85), and 9.76 (6.67), respectively. Most of the nurses were working in critical care wards and caring for critical patients. Other demographic data are presented in Table 1.

Mean of EI of nurses was 63.19 (8.22) and in different subscales of the EI self-awareness had the highest grade and self-management had the lowest grade. Details are summarized in Table 2.

Independent t -test did not show a statistically significant difference in the mean scores of emotional intelligence in terms of gender and type of ward. Also, the independent t -test showed that married nurses had higher EI scores compared to single nurses. The one-way ANOVA statistical test didn't show any significant difference in the scores of EI based on caring level, education, and employment status (Table 3).

Married nurses had a higher score in subscales of self-management and relationship management compared to singles. Also, nurses who were caring for more complex patients had a higher score in the relationship management subscale (Table 3).

Discussion

Based on available data the EI of nurses caring for COVID-19 patients is not well studied. This study was done to assess the EI of nurses caring for COVID-19 patients. Nurses had a moderate score in EI. The EI of married nurses was higher than single. In general, the highest and lowest scores were for self-awareness and self-management, respectively.

Different scores are reported for EI in different studies. Some studies report that the EI of nurses is optimal (Barkhordari et al., 2016; Konstantinou et al., 2017; Rostami et al., 2016; Saeid et al., 2013). In some studies nurses' EI score is reported low (Codier et al., 2008; Harper & Jones-Schenk, 2012). The reason for this discrepancy could be the assessment tool or the definition of EI. Despite the difference in assessment tools, the results of these studies affirm that the EI of nurses needs more considerations and some measures should be planned. The nature of the nursing profession and the diversity of situations they face require the nurses have high EI (Barkhordari et al., 2016). The EI, during

Table 1
General information of respondents.

| Characteristics | | Frequency (%) |
|-------------------|--------------|---------------|
| Gender | Male | 98 (46.4) |
| | Female | 111 (52.6) |
| Marriage | Single | 58 (27.5) |
| | Married | 151 (71.6) |
| University degree | BSC | 153 (72.5) |
| | MSc | 56 (26.5) |
| Ward | General | 74 (35.1) |
| | ICU | 130 (61.6) |
| Level of Care | Mild | 13 (6.2) |
| | Moderate | 83 (39.3) |
| | Sever | 110 (52.1) |
| Employment | Formal | 144 (68.2) |
| | Informal | 51 (24.2) |
| | Partial time | 10 (4.7) |

Most of nurses participated in this study were Female, married, and were working in ICU wards.

Table 2
Mean Score and standard deviation Emotional Intelligence and Components.

| Variable | Mean (SD) |
|-------------------------|---------------|
| Emotional intelligence | 63.19 (8.22) |
| Self-awareness | 68.83 (11.46) |
| Self-management | 56.41 (9.99) |
| Social awareness | 63.24 (10.39) |
| Relationship management | 66.55 (11.47) |

Totally, nurses' emotional intelligence was moderate. Also, self-awareness and self-management had highest and lowest scores, respectively.

situations such as the COVID-19 outbreak, could have a protective role for negative emotions such as fear, anxiety, and sadness (Morón & Biolik-Moroñ, 2020). This issue is more important due to the effects of a pandemic on the occurrence of psychological symptoms and the mental health of health care workers (Di Tella et al., 2020; Sampaio et al., 2020). One of the main challenges managers are faced is the psychological well-being of health care workers (Greenberg et al., 2020; Maben & Bridges, 2020).

This issue could have multiple negative consequences such as job dissatisfaction, stress, burnout, reduced performance, and the quality of care. EI gives the nurses the power to think better during critical and difficult situations and make better decisions by controlling the psychological reactions (Kelishami et al., 2017). Also, EI is an effective strategy for reducing job stress and promoting resiliency (Cleary et al., 2018; Khoshnazary et al., 2016). In terms of clinical performance, improving EI will lead to improving the quality of nursing care (Codier et al., 2013; Jang et al., 2016).

Self-awareness and self-management had the highest and lowest scores, respectively. This means that nurses have a high competency in recognizing their emotions, but their ability to manage them is not desirable. The results of similar studies also confirm the findings of the present study (Ko & Kim, 2014; Mosadegh Rad, 2015; Stiglic et al., 2018). Trying to improve the self-management is necessary. Self-management causes better performance in different stressful situations such as discordant with colleagues, denial, and emotions of patients and family members (Meires, 2018).

Considering the dimensions of EI is very important. Because we can assess the nurse and diagnose the area the person is weak and plan for improving it. Self-awareness and self-management are considered as individual capabilities of emotional intelligence and social awareness and relationship management as social capabilities (Ganji et al., 2006). Therefore, we should not focus only on the overall score of emotional intelligence, but the score of dimensions are more important.

Married nurses had higher scores in EI. Also, the score of self-management and relationship management were higher in married nurses. This difference also has been mentioned in other studies (Ebrahimi et al., 2020; Kalyoncu et al., 2012). This can be due to the involvement of married people in issues such as personal disputes and unfulfilled expectations that require empathy and adjustment. Therefore, it can be an exercise to improve and develop emotional intelligence. There is no significant relationship between gender and age with emotional intelligence in other similar studies (Khandan et al., 2015; Kooker et al., 2007; Petrovici & Dobrescu, 2014). However, differences in the type of tools used to assess emotional intelligence can also be particularly effective.

In the present study, there was no difference between the EI of nurses caring for patients with COVID19 in the normal and critical wards. In terms of dimensions of emotional intelligence, nurses caring for clinically stable COVID19 patients had a higher relationship management score. In the literature review, the results indicate a moderate and low level of emotional intelligence in nurses working in the intensive care unit (Delpasand et al., 2011; Saeed et al., 2013).

However, COVID19 pandemic affects the mental health of health

Table 3
The relationship between emotional intelligence and Dimensions with demographic characteristics.

| | Gender | | Marriage | | University degree | | Ward | | Level of Care | | | Employment | | Partial time |
|---|---|---|---|---|---|--|---|--|--|---|---|---|---|--|
| | Male | Female | Single | Married | BSC | MSc | General | ICU | Mild | Moderate | Severe | Formal | Informal | |
| Emotional intelligence (mean (SD)) Test | 63.43 (7.80) t-Test T = 0.51, P = 0.6 | 62.8 (8.55) t-Test T = 2.27, P = 0.02 | 61 (8.48) t-Test ^a T = 2.27, P = 0.02 | 64 (7.96) t-Test T = 1.18, P = 0.23 | 62.71 (8.05) t-Test T = 1.18, P = 0.23 | 64.22 (8.54) t-Test T = 1.36, P = 0.13 | 63.99 (8.27) t-Test T = 1.36, P = 0.13 | 62.38 (8.02) t-Test T = 0.17, P = 0.86 | 59.29 (8.07) ANOVA F = 2.49, P = 0.08 | 64.29 (8.12) ANOVA F = 2.49, P = 0.08 | 62.61 (8.24) ANOVA F = 2.49, P = 0.08 | 62.60 (7.02) ANOVA F = 0.15, P = 0.85 | 63.26 (7.02) ANOVA F = 0.15, P = 0.85 | 64.43 (8.63) ANOVA F = 0.15, P = 0.85 |
| Self-awareness (mean (SD)) Test | 68.54 (11.78) t-Test T = 0.22, P = 0.82 | 68.89 (11.23) t-Test T = 0.22, P = 0.82 | 66.26 (12.18) t-Test T = 1.93, P = 0.054 | 69.67 (11.07) t-Test T = 0.70, P = 0.48 | 68.39 (11.11) t-Test T = 0.70, P = 0.48 | 69.64 (12.4) t-Test T = 0.70, P = 0.48 | 68.65 (11.89) t-Test T = 0.17, P = 0.86 | 68.36 (11.3) t-Test T = 0.17, P = 0.86 | 68.21 (12.4) ANOVA F = 0.13, P = 0.87 | 69.08 (11.43) ANOVA F = 0.13, P = 0.87 | 68.24 (11.34) ANOVA F = 0.13, P = 0.87 | 68.54 (11.22) ANOVA F = 0.13, P = 0.87 | 67.58 (10.58) ANOVA F = 0.13, P = 0.87 | 69.00 (14.66) ANOVA F = 0.13, P = 0.87 |
| Self-management (mean (SD)) Test | 57.60 (9.48) t-Test T = 1.67, P = 0.9 | 55.30 (10.30) t-Test T = 1.67, P = 0.9 | 53.95 (9.49) t-Test ^a T = 2.20, P = 0.029 | 57.31 (10.02) t-Test T = 1.58, P = 0.11 | 55.72 (10.12) t-Test T = 1.58, P = 0.11 | 58.17 (9.38) t-Test T = 1.58, P = 0.11 | 56.82 (10.08) t-Test T = 0.58, P = 0.56 | 55.97 (9.97) t-Test T = 0.58, P = 0.56 | 50.43 (9.53) ANOVA F = 2.69, P = 0.07 | 57.32 (10.86) ANOVA F = 2.69, P = 0.07 | 56.59 (9.32) ANOVA F = 2.69, P = 0.07 | 55.99 (10.09) ANOVA F = 2.69, P = 0.07 | 56.82 (9.33) ANOVA F = 2.69, P = 0.07 | 56.67 (10.89) ANOVA F = 2.69, P = 0.07 |
| Social awareness (mean (SD)) Test | 63.06 (10.09) t-Test T = 0.10, P = 0.91 | 63.21 (10.6) t-Test T = 0.10, P = 0.91 | 63.17 (11.25) t-Test T = 0.02, P = 0.97 | 63.13 (10.05) t-Test T = 0.83, P = 0.40 | 63.50 (9.75) t-Test T = 0.83, P = 0.40 | 62.14 (11.9) t-Test T = 0.83, P = 0.40 | 64.11 (11.01) t-Test T = 1.09, P = 0.27 | 62.46 (9.89) t-Test T = 1.09, P = 0.27 | 62.77 (11.1) ANOVA F = 1.55, P = 0.22 | 64.53 (10.87) ANOVA F = 1.55, P = 0.22 | 61.93 (9.77) ANOVA F = 1.55, P = 0.22 | 62.67 (10.31) ANOVA F = 1.55, P = 0.22 | 63.61 (10.28) ANOVA F = 1.55, P = 0.22 | 62.80 (11.00) ANOVA F = 1.55, P = 0.22 |
| Relationship management (mean (SD)) Test | 66.38 (10.17) t-Test T = 0.11, P = 0.91 | 66.55 (12.58) t-Test T = 0.11, P = 0.91 | 63.62 (12.10) t-Test ^a T = 2.24, P = 0.026 | 67.57 (11.09) t-Test T = 1.36, P = 0.17 | 65.82 (11.21) t-Test T = 1.36, P = 0.17 | 68.26 (12.1) t-Test T = 1.36, P = 0.17 | 68.48 (11.61) t-Test T = 2.07, P = 0.03 | 65.06 (11.1) t-Test T = 2.07, P = 0.03 | 60.38 (9.83) ANOVA ^a F = 3.40, P = 0.03 | 68.40 (11.00) ANOVA ^a F = 3.40, P = 0.03 | 65.59 (11.73) ANOVA ^a F = 3.40, P = 0.03 | 65.54 (11.62) ANOVA ^a F = 3.40, P = 0.03 | 67.06 (10.79) ANOVA ^a F = 3.40, P = 0.03 | 70.75 (9.93) ANOVA ^a F = 3.40, P = 0.03 |

^a Married nurses had higher levels of emotional intelligence. In terms of dimensions of emotional intelligence, married nurses working in the general wards and caring for patients with moderate disease severity had a higher level of relationship management.

care workers and can affect the quality of care provided. Therefore, it is necessary to pay attention to the emotional intelligence of nurses caring for critically ill patients with COVID19.

Limitations

Due to limited access to research samples, questionnaires were sent and completed online. So we could not comply fully with our sampling schedule and plan.

Conclusion

The EI of nurses caring for patients with COVID19 was moderate. In terms of dimensions, the highest score was related to individual abilities (self-awareness and self-management). According to the results of the present study, the development of emotional intelligence based on the characteristics of nurses is very important in recognizing and managing their emotions. Due to the continuation of this pandemic and the possibility of mental and physical fatigue of health care workers, improving emotional intelligence can be effective in resilience and stability of the psychological status of employees.

Declaration of competing interest

We declare that we have no conflicts of interest.

Acknowledgment

The authors grateful to the Faculty of Nursing and trauma research center, Baqiyatallah University of Medical Sciences, for financially supporting this study and all the nurses who kindly cooperated in conducting this research.

References

- Alonazi, W. B. (2020). The impact of emotional intelligence on job performance during COVID-19 crisis: A cross-sectional analysis. *Psychology Research and Behavior Management*, 13, 749.
- Baba, M. M. (2020). Navigating COVID-19 with emotional intelligence. *International Journal of Social Psychiatry*, 66(8), 810–820.
- Barkhordari, M., Rostambeygi, P., & Ghasemnejad, M. (2016). Measuring emotional intelligence of nursing student. *Iranian Journal of Nursing Research*, 10(4), 12–19.
- Cleary, M., Visentin, D., West, S., Lopez, V., & Kornhaber, R. (2018). Promoting emotional intelligence and resilience in undergraduate nursing students: An integrative review. *Nurse Education Today*, 68, 112–120.
- Codier, E., Freitas, B., & Muneno, L. (2013). *Developing emotional intelligence ability in oncology nurses: A clinical rounds approach. Paper presented at the Oncology Nursing Forum.*
- Codier, E., Kooker, B. M., & Shultz, J. (2008). Measuring the emotional intelligence of clinical staff nurses: An approach for improving the clinical care environment. *Nursing Administration Quarterly*, 32(1), 8–14.
- Delpasand, M., Nasiripoor, A. A., Raiisi, P., & Shahabi, M. (2011). *Relationship between emotional intelligence and occupational burnout among nurses in Critical Care Units.*
- Di Tella, M., Romeo, A., Benfante, A., & Castelli, L. (2020). Mental health of healthcare workers during the COVID-19 pandemic in Italy. *Journal of Evaluation in Clinical Practice*, 26(6), 1583–1587.
- Doherty, E. M., Cronin, P. A., & Offiah, G. (2013). Emotional intelligence assessment in a graduate entry medical school curriculum. *BMC Medical Education*, 13(1), 38.
- Ebrahimi, N., Sadeghmoghadam, L., Hosseinzadeh, F., & Bahri, N. (2020). The relationship between emotional intelligence with job and individual characteristics of nursing staff. *Journal of Health and Safety at Work*, 10(3).
- Ganji, H., Mirhashemi, M., & Sabet, M. (2006). Bradberry-Greaves' emotional intelligence test: Preliminary norming-process. *Journal of Thought & Behavior*, 1(2), 23–35.
- Greenberg, N., Docherty, M., Gnanapragasam, S., & Wessely, S. (2020). Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*, 368.
- Harper, M. G., & Jones-Schenk, J. (2012). The emotional intelligence profile of successful staff nurses. *The Journal of Continuing Education in Nursing*, 43(8), 354–362.
- Huang, L., Lin, G., Tang, L., Yu, L., & Zhou, Z. (2020). Special attention to nurses' protection during the COVID-19 epidemic. *BioMed Central*, 24(120), 1–3.
- Jang, R.-j., Kang, Y.-S., & Kim, Y.-m. (2016). The relationships in emotional intelligence, job satisfaction, and quality of nursing service in hospital nurses. *Journal of the Korea Academia-Industrial Cooperation Society*, 17(6), 326–337.
- Kalyoncu, Z., Güney, S., Arslan, M., & Ayranci, E. (2012). *Analysis of the relationship between emotional intelligence and stress caused by the organization: A study of nurses.*
- Kelishami, F. G., Farahani, M. A., Orak, R. J., Ameri, Z. A., Hashemi, S. B., & Seyedfatemi, N. (2017). Emotional intelligence in nursing, models and methods of measurement. *Advances in Nursing & Midwifery*, 26(93), 21–29.
- Khandan, M., Kavousi, A., & Koohoei, A. (2015). Application of structural equations modeling to assess relationship among emotional intelligence, general health and occupational accidents. *International Journal of Occupational Hygiene*, 7(3), 124–131.
- Khoshnazary, S., Hosseini, M., Fallahi Khoshknab, M., & Bakhshi, E. (2016). The effect of Emotional Intelligence (EI) training on nurses' resiliency in Department of Psychiatry. *Iranian Journal of Psychiatric Nursing*, 3(4), 28–37.
- Ko, H. R., & Kim, J.-H. (2014). The relationships among emotional intelligence, interpersonal relationship, and job satisfaction of clinical nurses. *The Journal of Korean Academic Society of Nursing Education*, 20(3), 413–423.
- Konstantinou, M. S., Efstathiou, A., Charalambous, G., Kaitelidou, D., & Jelastopulu, E. (2017). Emotional intelligence and its impact on the emotional factors among nurses. *Journal of Education, Society and Behavioural Science*, 1–13.
- Kooker, B. M., Shultz, J., & Codier, E. E. (2007). Identifying emotional intelligence in professional nursing practice. *Journal of Professional Nursing*, 23(1), 30–36.
- Maben, J., & Bridges, J. (2020). Covid-19: Supporting nurses' psychological and mental health. *Journal of Clinical Nursing*, 29(15-16), 2742–2750.
- Meires, J. (2018). The essentials: Using emotional intelligence to curtail bullying in the workplace. *Urologic Nursing*, 38(3), 150–154.
- Mo, Y., Deng, L., Zhang, L., Lang, Q., Liao, C., Wang, N., ... Huang, H. (2020). Work stress among Chinese nurses to support Wuhan in fighting against COVID-19 epidemic. *Journal of Nursing Management*, 28(5), 1002–1009.
- Montes-Berges, B., & Augusto, J. M. (2007). Exploring the relationship between perceived emotional intelligence, coping, social support and mental health in nursing students. *Journal of Psychiatric and Mental Health Nursing*, 14(2), 163–171.
- Moroń, M., & Biolik-Moroń, M. (2020). Trait emotional intelligence and emotional experiences during the COVID-19 pandemic outbreak in Poland: A daily diary study. *Personality and Individual Differences*, 168, Article 110348.
- Moroń, M., Biolik-Moroń, M. J. P., & Differences, I. (2021). Trait emotional intelligence and emotional experiences during the COVID-19 pandemic outbreak in Poland: A daily diary study. 168, Article 110348.
- Morris, T., & Wakefield, S. (2008). *How do we ensure our curriculum has an impact on the Emotional Intelligence (EQ) of our learners. Paper presented at the 8th International Practice Development Conference.* Univ Northampton.
- Mosadegh Rad, A. M. (2015). The relationship of emotional intelligence and its dimensions on communication skills among emergency unit nurses. *Journal of Hospital*, 14(3), 67–74.
- Nightingale, S., Spiby, H., Sheen, K., & Slade, P. (2018). The impact of emotional intelligence in health care professionals on caring behaviour towards patients in clinical and long-term care settings: Findings from an integrative review. *International Journal of Nursing Studies*, 80, 106–117.
- Petrovici, A., & Dobrescu, T. (2014). The role of emotional intelligence in building interpersonal communication skills. *Procedia-Social and Behavioral Sciences*, 116, 1405–1410.
- Prieseemann, V., Balling, R., Brinkmann, M. M., Ciesek, S., Czypionka, T., Eckerle, L., ... Hotulainen, P. J. T. L. (2021). An action plan for pan-European defence against new SARS-CoV-2 variants. 397(10273), 469–470.
- Rostami, M., Movaghari, M., Taghavi, T., & Mehran, A. (2016). The relationship between emotional intelligence and coping styles of nurses in hospitals in Kermanshah University of Medical Sciences. *Iranian Journal of Nursing Research*, 11(1), 51–61.
- Saeed, Y., Javadi, M., & MOKHTARI, N. J. (2013). Assessment and comparison of emotional intelligence of nurses in general and intensive care units. *IJCCN*, 5(4), 214–221.
- Saeid, Y., Javadi, M., Mokhtari Noori, J., & Sirati Nir, M. (2013). On the relationship between emotional intelligence and demographical variables in nurses. *Journal of Military Medicine*, 15(1), 87–92.
- Sampaio, F., Sequeira, C., & Teixeira, L. (2020). Nurses' mental health during the Covid-19 outbreak: A cross-sectional study. *Journal of Occupational and Environmental Medicine*, 62(10), 783–787.
- Smith, G. D., Ng, F., & Li, W. H. C. (2020). COVID-19: Emerging compassion, courage and resilience in the face of misinformation and adversity. *Journal of Clinical Nursing*, 29(9–10), 1425.
- Sommer, S. A., Howell, J. M., & Hadley, C. N. (2016). Keeping positive and building strength: The role of affect and team leadership in developing resilience during an organizational crisis. *Group & Organization Management*, 41(2), 172–202.
- Štiglic, G., Cilar, L., Novak, Z., Vrbnjak, D., Stenhouse, R., Snowden, A., & Pajnkihar, M. (2018). Emotional intelligence among nursing students: Findings from a cross-sectional study. *Nurse Education Today*, 66, 33–38.
- Talman, K., Hupli, M., Rankin, R., Engblom, J., & Haavisto, E. (2020). Emotional intelligence of nursing applicants and factors related to it: A cross-sectional study. *Nurse Education Today*, 85, Article 104271.
- Wu, W., Zhang, Y., Wang, P., Zhang, L., Wang, G., Lei, G., ... Xie, S. (2020). Psychological stress of medical staffs during outbreak of COVID-19 and adjustment strategy. *Journal of Medical Virology*, 92(10), 1962–1970.